

Vol. 101, March 2006



***Arabidopsis PHT1* promoter available for root-specific expression of heterologous genes in dicots and monocots.**

The promoter of *PHT1* encoding phosphate transporter of *Arabidopsis thaliana* was fused to β -glucuronidase (*GUS*) gene and the construct (*PHT1* promoter::*GUS*) was introduced into *Arabidopsis* and rice. Histochemical localization of *GUS* activity of a T1 rice plant indicated that plantlet exhibits strong *GUS* activity in roots (above left), *GUS* signal is not found in the leaves (above right), strong *GUS* signal is observed in the roots except for root tip (below left), *GUS* signaling is stronger in cells that generate root hairs than in cells that do not (below middle), and that root hair cells show strong *GUS* activity (transverse section).

The results obtained indicated that the dicot promoter can function efficiently in monocot plants, and that the *PHT1* promoter is a practical promoter for root-specific expression of heterologous genes both in dicots and monocots.

Related article: Koyama, T., Ono, T., Shimizu, M., Jinbo, T., Mizuno, R., Tomita, K., Mitsukawa, N., Kawazu, T.,

Kimura, T., Ohmiya, K., and Sakka, K., "**Promoter of *Arabidopsis thaliana* phosphate transporter gene drives root-specific expression of transgene in rice**", **J. Biosci. Bioeng., vol.99, 38-42 (2005)**.

[⇒JBB Archive Top](#)

[⇒JBB Archive: Vol. 93 \(2002\)–Vol. 106 \(2008\)](#)