

Vol. 106, September 2008



**Enrichment of carotenoids in flaxseed.**

Left: an untransformed flax plant (WARD cultivar). The transgenic flax plants were the same to untransformed control plants in their appearance except for seed color.

Right top: section of an untransformed flaxseed.

Right below three: sections of transgenic flaxseeds. These transgenic plants were generated by introduction of the phytoene synthase gene (*crtB*) derived from soil bacterium *Pantoea ananatis* (formerly called *Erwinia uredovora* 20D3).

The inner color of the transgenic flaxseeds was altered into orange due to the accumulation of  $\beta$ -carotene and  $\alpha$ -carotene. Total carotenoid amounts in these seeds were 65.4-156.3  $\mu\text{g/g}$  fresh weight, which corresponded to 7.8- to 18.6-fold increase, compared with those of untransformed controls.

Related article: Fujisawa, M., Watanabe, M., Choi, S.-K., Teramoto, M., Ohyama, K., and Misawa, N., "Enrichment of carotenoids in flaxseed (*Linum usitatissimum*) by metabolic engineering with introduction of bacterial phytoene synthase gene *crtB*" *J. Biosci. Bioeng.*, vol. 105, 636-641 (2008).

⇒JBBアーカイブ : Vol.107 (2009) ~最新号

⇒JBBアーカイブ : Vol. 93 (2002) ~Vol. 106 (2008)